








Figure 1



2/12

Figure 2A: Body weight change over study.




Figure 2B: Food intake over study.




Figure 2C: Neurological/behavioural score over study.







Figure 2D: Representative histological striatal sections of A) sham-operated rat, B) untreated rat, and C) PMX53-treated rat.



Sham (Day 14)
No cell loss



Untreated (Day 14)
Cell necrosis



PMX53-treated (Day
14)
Mild cell necrosis

4/12

Figure 3A: Body weight change after 7 days.




Figure 3B: Food intake after 7 days.





Figure 3C: Neurological/behavioural score over study.



5/12


Figure 3D: Representative histological striatal sections of A) sham-operated rat, B) untreated rat, C) PMX53-treated rat and, D) PMX205-treated rat.



Sham (Day 7)
No cell loss



Untreated (Day 7)
Major cell necrosis



PMX53-treated (Day 7)
Minor cell necrosis



PMX205-treated (Day 7)
Minor cell necrosis

6/12

Figure 4A: Body weight change after 7 days.





Figure 4B: Food intake after 7 days.



7/12

Figure 4C: Neurological/behavioural score after 7 days.




Figure 4D: Striatal lesion size after 7 days.






Figure 5a




Figure 5b



Sham. 40x



Untreated.




9/12


Figure 5c



Sham. 40x



Untreated. 20x




Untreated. 40x



PMX205-treated.

10/12

Figure 6a



Untreated. 40x

Figure 6b



Untreated. 200x

11/12

Figure 7a




Figure 7b




Figure 7c





Figure 7d



12/12

Figure 8

